

RECKON 280SL Herbicide

A non-selective herbicide for post emergence broadcast use on canola, corn, cotton and soybean designated as LibertyLink®. Reckon 280SL may be used for weed control in non-LibertyLink® cotton when applied with a hooded sprayer. Reckon 280SL may also be applied as a broadcast burndown application before planting or prior to emergence of any conventional or transgenic variety of canola, corn, sweet corn*, cotton, olive, rice*, soybean or sugar beet. Reckon 280SL may be used for post emergence weed control in listed tree, vine and berry crops and rice*. Reckon 280SL may also be applied for potato vine desiccation.

*Not for use in CA

ACTIVE INGREDIENT:

Glufosinate-ammonium (CAS No. 77182-82-2).....	24.5%**
OTHER INGREDIENTS.....	<u>75.5%</u>
TOTAL	100.0%

**Equivalent to 2.34 pounds of active ingredient per U.S. gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Inside Booklet. **For MEDICAL emergencies call the National Poison Control Center at 1-800-222-1222 and for TRANSPORTATION emergencies call Chemtrec at 1-800-424-9300.**

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
--------------	--

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. For MEDICAL emergencies call the National Poison Control Center at 1-800-222-1222.

NOTE TO PHYSICIAN

If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.



**Manufactured for:
Solera ATO, LLC
12230 E Del Norte
Yuma, AZ 85367**

**EPA Reg. No.: 88685-2-84237
EPA Est. No.: 75486-CHN-001
Net Contents: 2.5gal
03-RKN-R030716**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants;
- shoes and socks; and
- waterproof or chemical-resistant gloves.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When mixing and loading wear a chemical-resistant apron. For overhead exposure wear chemical-resistant headgear. When cleaning equipment wear a chemical-resistant apron.

Engineering Control Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present. Do not apply to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions, this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur to minimize water runoff is recommended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In the State of New York Only: Not For Use in Nassau and Suffolk Counties.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 12 hours with the exception of sweet corn irrigation activities which have a 4 day REI. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls worn over short-sleeved shirt and short pants; chemical resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils; chemical resistant footwear plus socks; protective eyewear (goggles, face shield or safety glasses).

IMPORTANT CROP SAFETY INFORMATION READ BEFORE USING THIS PRODUCT

Reckon 280SL may be applied as a **burndown treatment prior to planting or prior to emergence** of any conventional or transgenic variety of canola, corn, sweet corn*, cotton, olive, rice*, soybean or sugar beet. Reckon 280SL may be applied to conventional cotton not tolerant to the active ingredient in Reckon 280SL using a hooded sprayer.

*Not for use in CA

Post emergence row crop applications of Reckon 280SL may be made only to crops tolerant to the active ingredient in this product. Orion GFS, LLC does not warrant the use of this product on crops other than those designated as LibertyLink® to safely withstand the application of Reckon 280SL.

The basis of selectivity of Reckon 280SL in crops is the presence of a gene in LibertyLink® crops which results in a plant that is tolerant to the active ingredient in Reckon 280SL. Crops not containing this gene will not be tolerant to Reckon 280SL, and severe crop injury and/or death may occur. Do not allow spray to contact foliage or green tissue of desirable vegetation other than crops tolerant to the active ingredient in this product.

Applications to trees, vines and berries should avoid contact of Reckon 280SL solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees, berries and vines. Only trunks with callused, mature brown bark should be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of Reckon 280SL with parts of trees, berries or vines other than mature brown bark can result in serious damage.

INFORMATION

Reckon 280SL is a water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds in LibertyLink® canola, corn, cotton and soybean, and in trees, vines and berries. Reckon 280SL may also be applied for potato vine desiccation. Reckon 280SL may also be applied as a broadcast burndown application before planting or prior to emergence of any conventional or transgenic variety of canola, corn, sweet corn*, cotton, olive, rice*, soybean or sugar beet.

*Not for use in CA

Reckon 280SL is only foliar-active with little or no activity in soil. Weeds that emerge after application will not be controlled. Apply Reckon 280SL to actively growing weeds as described in the Weed Control Recommendations for Row Crops section to get maximum weed control. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Necrosis of leaves and young shoots occur within 2 to 4 days after application under good growing conditions.

- Reckon 280SL is rainfast four (4) hours after application to most weed species; therefore, rainfall within four (4) hours may necessitate retreatment or may result in reduced weed control.
- Applications should be made between dawn and 2 hours before sunset to avoid the possibility of reduced lambsquarters and velvetleaf control.

Consult your local Cooperative Extension Service or Orion GFS Representative for guidelines on the optimum application timing for Reckon 280SL in your region.

- Weed control may be reduced if application is made when heavy dew, fog and mist/rain are present; or when weeds are under stress due to environmental conditions such as drought, cool temperatures or extended periods of cloudiness.
- To maximize weed control, do not cultivate from 5 days before an application to 7 days after an application.

ROTATIONAL CROP RESTRICTIONS*

Rotational crop planting intervals following application of Reckon 280SL are listed below. Failure to comply with these restrictions may result in illegal residues in rotated crops.

Rotational Crop	Plant Back Interval (Minimum Rotational Crop Planting Interval from Last Application)
Canola, Corn, Sweet Corn, Cotton, Rice, Soybean and Sugar Beet	May be planted at any time
Root and Tuber Vegetables, Leafy Vegetables, Brassica Leafy Vegetables and Small Grains (barley, buckwheat, oats, rye, teosinte, triticale, and wheat)	70 Days
All Other Crops	180 Days

*See Application Directions for Potato Vine Desiccation for Rotational Crop Restrictions specifically after Reckon 280SL applications to potatoes.

Integrated Weed Management

The active ingredient in Reckon 280SL is glufosinate-ammonium, which is a glutamine synthetase inhibitor (Group 10). Integrated weed management guidelines promote an economically viable, environmentally sustainable and socially acceptable weed control program regardless of the herbicide(s) used. The highlights of a successful integrated weed management program include:

1. Correctly identify weeds and look for trouble areas within field to identify resistance indicators.
2. Rotate crops.
3. Start the year with clean fields.
4. Rotate herbicide modes of action by using multiple modes of action during the year and apply no more than two applications of a single herbicide mode of action to the same field in a two-year period. One method to accomplish this is to rotate herbicide tolerant trait systems.
5. Apply listed rates of herbicides to actively growing weeds at the correct time with the right application techniques.
6. Control any weeds that may have escaped the herbicide application.
7. Thoroughly clean field equipment between fields.

Contact your local agronomic advisor for more specific information on integrated weed management for your area.

WEED CONTROL FOR ROW CROPS

Rates in ounces of formulated product per acre for the control of weeds at selected heights are shown in the weed control tables. In weed populations with mixed species, apply at a rate needed for the species that requires the highest rate.

Broadleaf Weed Control					
Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	22 fl oz/A	29 fl oz/A ^{ab}		22 fl oz/A	29 fl oz/A ^{ab}
Amaranth, Palmer ³	NR	4	Morningglory, sharppod ³	2	4
Anoda, spurred	3	5	Morningglory, smallflower ³	4	6
Beggarweed, Florida	4	5	Morningglory, tall ³	6	8
Black medic	5	7	Mustard, wild	4	6
Blueweed, Texas	5	7	Nightshade, black	4	6
Buckwheat, wild	6	7	Nightshade, eastern black	6	8
Buffalobur	6	7	Nightshade, hairy	6	8
Burcucumber	6	10	Pennycress (stinkweed)	4	6
Catchweed bedstraw (cleavers)	2	4	Pigweed, redroot ³	3	4
Carpetweed	4	6	Pigweed, prostrate ³	3	4
Chickweed, common	6	8	Pigweed, spiny ³	3	4
Cocklebur, common	6	14	Pigweed, smooth ³	3	4
Copperleaf, hophornbeam	4	6	Pigweed, tumble ³	3	4
Cotton, volunteer ¹	6	8	Puncturevine	4	6
Croton, tropic	3	5	Purslane, common	2	4
Croton, woolly	2	4	Pusley, Florida	S	3
Eclipta	4	6	Ragweed, common	6	10
Devil's claw	2	4	Ragweed, giant	6	12
Fleabane, annual	6	8	Senna coffee	4	6
Galinsoga, hairy	6	8	Sesbania, hemp	6	8
Galinsoga, smallflower	6	7	Shepherd's-purse	6	8
Groundcherry, cutleaf	4	5	Sicklepod (java bean)	4	6
Geranium, cutleaf	4	6	Sida, prickly	4	5
Hempnettle	4	6	Smartweed, Pennsylvania	6	14
Horsenettle, Carolina ²	2	4	Smellmelon	4	6
Jimsonweed	6	10	Sowthistle, annual	6	8
Knotweed	3	5	Soybeans, volunteer ¹	6	8
Kochia	4	6	Spurge, prostrate	2	4
Ladysthumb	6	14	Spurge, spotted	2	4
Lambsquarters, common ³	4	6	Starbur, bristly	4	6
Mallow, common	4	6	Sunflower, common	6	14
Mallow, Venice	6	8	Sunflower, prairie	3	5
Marestail	S	6-12	Sunflower, volunteer	6	10
Marshelder, annual	4	6	Thistle, Russian ²	S	6-12
Morningglory, entireleaf ³	6	8	Velvetleaf	3	4
Morningglory, ivyleaf ³	6	8	Waterhemp, common ³	NR	5
Morningglory, pitted ³	6	8	Waterhemp, tall ³	NR	5

^aIn cotton, Reckon 280SL may be applied at 29 fl oz/A three times per year.

^bDo not apply more than 22 fl oz/A of Reckon 280SL post emergence in a single application to canola and corn.

S Indicates suppression

¹ Volunteer LibertyLink[®] crops from the previous season will not be controlled.

² May require sequential applications for control.

³ For applications to corn, tank mixing with atrazine may enhance weed control of this species.

NR Not Recommended

Grass Weed Control					
Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	22 fl oz/A	29 fl oz/A ^{ab}		22 fl oz/A	29 fl oz/A ^{ab}
Barley, volunteer ³	3	4	Millet, wild proso	6	7
Barnyardgrass	3	5	Millet, proso volunteer	6	7
Bluegrass, annual	3	5	Oat, wild ²	3	4
Corn, volunteer ¹	10	12	Panicum, fall	3	5
Crabgrass, large ²	3	5	Panicum, Texas	4	6
Crabgrass, smooth ²	3	5	Rice, red	4	6
Cupgrass, woolly	6	12	Rice, volunteer ¹	4	6
Foxtail, bristly	6	8	Sandbur, field ²	S	2
Foxtail, giant	6	12	Shattercane	6	8
Foxtail, green	6	12	Signalgrass, broadleaf	3	5
Foxtail, robust purple	6	8	Sprangletop	4	6
Foxtail, yellow ²	3	4	Sorghum, volunteer	6	8
Goosegrass ³	2	3	Stinkgrass	4	6
Johnsongrass, seedling	3	5	Wheat, volunteer ²	4	5
Junglerice	3	5	Witchgrass	4	6

^a In cotton, Reckon 280SL may be applied at 29 fl oz/A three times per year.

^b Do not apply more than 22 fl oz/A of Reckon 280SL post emergence in a single application to canola and corn.

S Indicates suppression

¹ Volunteer LibertyLink[®] crops from the previous season will not be controlled. A timely cultivation 7 to 10 days after an application and/or retreatment 10-21 days after the first application for controlling dense clumps of volunteer LibertyLink[®] crops.

² For best control of yellow foxtail, field sandbur, crabgrass, and wild oats, treat prior to tiller initiation.

³ A sequential application may be necessary for control.

Biennial and Perennial Weeds			
For control of the biennial and perennial weeds listed below, apply tank mix partners or sequential applications of Reckon 280SL (22 fl oz/A followed by 22 fl oz/A).			
Alfalfa	Burdock	Goldenrod, gray*	Orchardgrass
Artichoke, Jerusalem	Bursage, woolyleaf	Johnsongrass, rhizome	Poinsettia, wild
Bermudagrass	Chickweed, mouse-ear	Milkweed, common*	Pokeweed
Bindweed, field	Clover, Alsike	Milkweed, honeyvine*	Quackgrass*
Bindweed, hedge	Clover, red	Muhly, wirestem*	Sowthistle, perennial
Bluegrass, Kentucky	Dandelion	Nightshade, silverleaf	Thistle, bull
Blueweed, Texas	Dock, smooth	Nutsedge, purple*	Thistle, Canada
Bromegrass, smooth	Dogbane, hemp*	Nutsedge, yellow*	Timothy*
			Wormwood, biennial

*Suppression Only

**See the *Application Directions for Use on Cotton* section of this label for additional use rates.

APPLICATION AND MIXING PROCEDURES

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment. Uniform thorough spray coverage is important to achieve consistent weed control.

Ground Application: Refer to the Rate Tables for proper application rates. DO NOT apply when winds are gusty or when conditions will favor movement of spray particles off the desired spray target. To avoid drift and insure consistent weed control, apply Reckon 280SL with the spray boom as low as possible while maintaining a uniform spray pattern. Reckon 280SL should be applied broadcast in a minimum of 15 gallons of water per acre using a minimum spray pressure of 40 psi and a maximum ground speed of 10 mph. The use of 80 degree or 110 degree flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration. Application of the spray at a 45-degree angle forward will result in better spray coverage. Under dense weed/crop canopies, 20 to 40 gallons of water per acre should be used so that thorough spray coverage will be obtained. Apply Reckon 280SL using nozzles and pressures that generate MEDIUM (about 250 to 350 microns) spray droplets category as reported by the nozzle manufacturer and in accordance to ASABE S 572. Do not use nozzles and pressures that result in COARSE sprays. FINE sprays should also be avoided to minimize spray drift risk. Boom height should be based on nozzle manufacturer recommendations. See the Spray Drift Management section of this label for additional information on proper application of Reckon 280SL.

Aerial Application: Poor coverage will result in reduced weed control. For optimal weed control, apply Reckon 280SL in a minimum of 10 gallons per acre. Apply Reckon 280SL using nozzles and pressures that generate MEDIUM (about 300 to 400 microns) spray droplets category as reported by the nozzle manufacturer and in accordance to ASABE S 572 based upon the selected air speed. Do not use nozzles and pressures that result in COARSE sprays. FINE sprays should also be avoided to minimize spray drift risk. See the Spray Drift Management section of this label for additional information on proper application of Reckon 280SL.

COMPATIBILITY TESTING

If Reckon 280SL is to be mixed with pesticide products not listed on this label, test the compatibility of the intended tank mixture prior to mixing the products in the spray tank. The following procedure assumes a spray volume of 25 gallons per acre. For other spray volumes, adjust the amount of the water used accordingly. Check compatibility as follows:

1. Place 1.0 pint of water from the source that will be used to prepare the spray solution in a clear 1-quart jar.
2. For each pound of a dry tank mix partner to be applied per acre, add 1.5 teaspoons to the jar.
3. For each 16 fl oz of a liquid tank mix partner to be applied per acre, add 0.5 teaspoon to the jar.
4. For each 16 fl oz of Reckon 280SL to be applied per acre, add 0.5 teaspoon to the jar.
5. After adding all the ingredients, place a lid on the jar and tighten. Invert 10 times to mix.
6. Let the mixture stand for 15 minutes, and evaluate the solution for uniformity and stability. Look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. If the tank mix partners are not compatible, do not use the mixture in a spray tank.
7. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the Storage and Disposal section of this label.

MIXING INSTRUCTIONS

Tank Mix: Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing. Refer to the specific crop section for rate recommendations and other restrictions.

Reckon 280SL must be applied with properly calibrated and clean equipment. Reckon 280SL is formulated to mix readily in water. Prior to adding Reckon 280SL to the spray tank, ensure that the spray tank is thoroughly clean, particularly if a herbicide with the potential to injure crops was previously used (see Cleaning Instructions).

Mix Reckon 280SL with water to make a finished spray solution as follows:

1. Fill the spray tank half full with water.
2. Start agitation.
3. If mixing with a flowable/wettable powder tank mix partner. Prepare a slurry of the proper amount of the product in a small amount of water. Add the slurry to the spray tank.
4. Add the appropriate amount of ammonium sulfate (AMS) to the spray tank.
5. If mixing with a liquid tank mix partner, add the liquid mix partner next.
6. Complete filling the spray tank with water.
7. Add the proper amount of Reckon 280SL and continue agitation.
8. If foaming occurs, use a silicone-based antifoam agent.

Ensure that all spray system lines including pipes, booms, etc. have the correct concentration of spray solution by flushing out the spray system lines before starting the crop application.

If tank mix partners recommended on this label are added, maintain good agitation at all times until contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzles or line strainers must be 50 mesh or larger.

CLEANING INSTRUCTIONS

Before using Reckon 280SL, thoroughly clean bulk storage tank, refillable tank, nurse tanks, spray tank, lines, and filter, particularly if a herbicide with the potential to injure crops was previously used. Equipment should be thoroughly rinsed using a commercial tank cleaner.

After using Reckon 280SL, triple rinse the spray equipment and clean with a commercial tank cleaner before using for crops not labeled LibertyLink®. Make sure any rinsate or foam is thoroughly removed from spray tank and boom. Rinsate may be disposed following the pesticide disposal directions on this label.

SPRAY DRIFT MANAGEMENT

Spray drift may result in injury to non target crops or vegetation. To avoid spray drift, do not apply when wind speed is greater than 10 MPH or during periods of temperature inversions. Do not apply when weather conditions, wind speed, or wind direction may cause spray drift to non-target areas.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For all non-aerial applications, wind speed must be measured adjacent to the application site, on the upwind side, immediately prior to application.

Sensitive Areas: The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Do not apply under circumstances where possible drift to unprotected persons or to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.

Aerial Drift Management: The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions on next page). **AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.**

Controlling Droplet Size:

Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure: Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles: Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation: Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind: Drift potential is lowest between wind speeds of 2 -10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Applications should be avoided below 2 miles per hour due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry. Avoid spraying during conditions of low humidity and/or high temperatures.

Temperature Inversions: Do not make aerial or ground applications into areas of temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

APPLICATION DIRECTIONS FOR BURNDOWN USE

Reckon 280SL may be applied as a **burndown treatment prior to planting or prior to emergence** of any conventional or transgenic variety of canola, corn, sweet corn, cotton, rice, soybean, or sugar beet. Apply a minimum of 29 fl oz/A of Reckon 280SL for burndown of existing weeds just prior to planting or prior to emergence of canola, corn, sweet corn, cotton, rice, soybean, or sugar beet. For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures.

In cotton, if environmental conditions prevent timely application, a single application may be made of up to 43 fl oz/A of Reckon 280SL. **If more than 29 fl oz/A are used in any single application, the yearly total may not exceed 72 fl oz/A, including all application timings.**

In soybean, if environmental conditions prevent timely applications, a single application may be made of up to 36 oz/A of Reckon 280SL. **If 29-36 fl oz/A are used in a single burndown application, one additional in-season application may be made at up to 29 fl oz/A. The yearly total may not exceed 65 fl oz/A, including all application timings.**

In canola, corn, rice, and sugar beet, if environmental conditions prevent timely applications, a single application may be made of up to 36 fl oz/A. **No additional applications of Reckon 280SL may be made post emergence to the crop during the year.**

	Burndown	In-Season Applications (LibertyLink® Varieties Only)	Yearly Maximum
Cotton Use Pattern 1	29 fl oz/A	2 applications at 22-29 fl oz/A*	87 fl oz/A
Cotton Use Pattern 2	30-43 fl oz/A	1 application at 22-29 fl oz/A*	72 fl oz/A
Soybean Use Pattern	29-36 fl oz/A	1 application at 22-29 fl oz/A**	65 fl oz/A
Canola, Corn, Rice, Sugar Beet	29-36 fl oz/A	None	36 fl oz/A

* LibertyLink® cotton OR with hooded sprayer for non-LibertyLink varieties (see Cotton use directions)

** LibertyLink® soybean only (See Soybean use directions)

APPLICATION DIRECTIONS FOR USE ON SUGAR BEET

THOROUGH SPRAY COVERAGE IS VERY IMPORTANT. Reckon 280SL works best when weeds are actively growing. A cultivation may be made at least 5 days before a Reckon 280SL application or 5 days after a Reckon 280SL application.

APPLICATION TIMING

Applications of Reckon 280SL on sugar beet may be made from the cotyledon stage up to the 10-leaf stage of the sugar beet. Reckon 280SL is a foliar-active material with no soil-residual activity. For best results, apply to emerged, young actively growing weeds. Weeds that emerge after application will not be controlled. Reckon 280SL will have an effect on weeds that are larger than the recommended leaf stage; however, speed of activity and control may be reduced. Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness. Reckon 280SL is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best weed control and sugar beet yield, Reckon 280SL applications should begin when weeds are up to 1 inch in height or diameter. Repeat applications should be made when newly germinated weeds again reach 1 inch in height or diameter. Refer to the Rate Tables for Weed Control In Sugar Beets for selection of the proper rate dependent upon the weed species present and size.

A repeat application of Reckon 280SL or a tank mix application with a residual herbicide selected from the tank mix partners listed on this label will be needed to control weeds that have not yet emerged at the time of application.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON SUGAR BEETS

1. **DO NOT** apply more than 30 fl oz/A of Reckon 280SL in one application and **DO NOT** apply more than 60 fl oz/A of Reckon 280SL on the sugar beet crop per year.
2. **DO NOT** apply Reckon 280SL within 60 days of harvesting sugar beets.
3. **DO NOT** plant rotation crops in a field treated with Reckon 280SL within 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product. Corn, soybeans, canola, and sugar beets tolerant to the active ingredient of Reckon 280SL may be planted at any time.
4. **DO NOT** graze the treated crop or cut for hay.
5. **DO NOT** add surfactants. Anti-foams or drift control agents may be added if needed.
6. **DO NOT** apply Reckon 280SL if sugar beets show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
7. **DO NOT** apply this product through any type of irrigation system.

RATE TABLES FOR WEED CONTROL IN SUGAR BEETS

The rates of Reckon 280SL in fluid ounces (pints) of formulated product per acre to be used for the control of weeds at selected heights are shown in the following tables. In weed populations with mixed species, apply the rate needed for all species present.

Grass Weeds Controlled with Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Height)		Comments on Weed Growth Stage/ Application Timing/ Number of Applications
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)	
Barley, volunteer	1 - 2 leaf (2")	3 leaf (3")	Multiple applications may be required
Barnyardgrass	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Corn, volunteer	1 - 2 leaf (3")	3 - 4 leaf (6")	—
Crabgrass, large	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Crabgrass, smooth	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Cupgrass, woolly	1 - 5 leaf (4")	(8")	—
Foxtail, giant	1 - 4 leaf (3")	5 - 6 leaf (4")	Maximum of 2 tillers
Foxtail, green	1 - 4 leaf (3")	5 - 6 leaf (4")	Maximum of 2 tillers
Foxtail, yellow	1 - 3 leaf (2")	4 leaf (2")	Apply prior to tillering
Millet, volunteer proso	1 - 3 leaf (1")	4 - 5 leaf (3")	Maximum of 1 tiller
Millet, wild proso	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Oat, wild	1 - 2 leaf (2")	3 leaf (3")	Maximum of 1 tiller
Panicum, fall	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Panicum, Texas	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Sandbur, field	-	1 - 4 leaf (2")	Apply prior to tillering
Wheat, volunteer	1 - 2 leaf (2")	3 leaf (3")	Maximum of 1 tiller

* Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.

For improved control of heavy populations or larger than recommended volunteer wheat, volunteer barley, yellow foxtail, and wild oats, Reckon 280SL can be tank mixed with Assure® II Herbicide, Poast® Herbicide, Prism* Herbicide or Select® 2EC Herbicide.

Perennial Weeds Controlled by Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Height/Diameter)		Comments on Number of Applications
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)	
Quackgrass	-	1 - 3 leaf (3")	Multiple applications required
Sowthistle, perennial	-	1 - 4 leaf (3")	Multiple applications required
Thistle, Canada	-	1 - 4 leaf (3")	Multiple applications required

* Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.

Broadleaf Weeds Controlled by Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Diameter)	
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)
Buckwheat, wild	1 - 4 leaf (2")	5 - 6 leaf (3")
Buffalobur	1 - 4 leaf (2")	5 - 6 leaf (3")
Carpetweed	—	1 - 4 leaf (2")
Chickweed, common	1 - 4 leaf (2")	5 - 6 leaf (3")
Cocklebur, common	1 - 6 leaf (3")	7 - 8 leaf (5")
Kochia	(1")	(2")
Ladysthumb	1 - 2 leaf (1")	3 - 4 leaf (3")
Lambsquarter, common	1 - 2 leaf (1")	4 - 5 leaf (3")
Mallow, Venice	1 - 4 leaf (2")	5 - 6 leaf (3")
Marshelder	1 - 2 leaf (1")	3 - 4 leaf (2")
Mustard, wild	1 - 4 leaf (2")	5 - 6 leaf (3")
Nightshade, eastern black	1 - 4 leaf (2")	5 - 6 leaf (3")
Pigweed, prostrate	(1")	(3")
Pigweed, redroot	1 - 2 leaf (1")	3 - 4 leaf (3")
Pigweed, smooth	1 - 2 leaf (1")	3 - 4 leaf (3")
Pigweed, spiny	1 - 2 leaf (1")	3 - 4 leaf (3")
Purslane, common	(1")	(2")
Ragweed, common	1 - 6 leaf (3")	7 - 8 leaf (5")
Ragweed, giant	1 - 4 leaf (2")	5 - 6 leaf (3")
Shepherd's purse	1 - 4 leaf (2")	5 - 6 leaf (3")
Smartweed, Pennsylvania	1 - 2 leaf (1")	3 - 4 leaf (3")
Sowthistle, annual	1 - 4 leaf (2")	5 - 6 leaf (3")
Sunflower, common	1 - 6 leaf (3")	7 - 8 leaf (5")
Thistle, Russian	(1")	(2")
Velvetleaf	1 - 2 leaf (1")	3 - 4 leaf (3")

*Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.

APPLICATION DIRECTIONS FOR USE ON CANOLA

Apply Reckon 280SL only to canola labeled as LibertyLink®. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION RATE AND TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimal yield, early season weed removal is important.

Applications of Reckon 280SL on canola may be made from the cotyledon stage up to the early bolting stage of the canola. Slight discoloration of the canola may be visible after application. This effect is temporary and will not influence crop growth, maturity, or yield. Apply Reckon 280SL at 22 fl oz/A per application. A second application of Reckon 280SL may be needed to control weeds that have not yet emerged at the time of application.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON CANOLA

- **DO NOT** use on canola in the states of Alabama, Delaware, Georgia, Kentucky, Maryland, New Jersey, North Carolina, South Carolina, Tennessee, Virginia and West Virginia.
- **DO NOT** apply more than two applications of Reckon 280SL per year. Sequential applications should be at least 10 days apart.
- **DO NOT** apply Reckon 280SL within 65 days of harvesting canola.
- **DO NOT** apply more than 44 fl oz/A of Reckon 280SL per year. If Reckon 280SL was used in a burndown application, no post emergence applications may be applied to the crop.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply Reckon 280SL if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the "**Rotational Crop Restrictions**" section under the "**Information**" heading of this label for the appropriate rotational crop plant back intervals.

SPRAY ADDITIVES

Reckon 280SL must be applied with ammonium sulfate (AMS). Use only fine feed grade or spray grade AMS at 3 pounds per acre. Anti-foams or drift control agents may be added if needed. Use of additional surfactants or crop oils may increase risk of crop response.

CANOLA TANK MIX INSTRUCTIONS

Reckon 280SL at 22 fl oz/A plus AMS may be used in tank-mix combination with certain herbicides for improved control of larger than labeled grasses. Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the canola to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing. The AMS rate may be reduced to 1.5 lb/A when Reckon 280SL is tank mixed with a reduced rate of one of the grass herbicides specified below.

TANKMIX PARTNERS FOR RECKON 280SL ON INVIGOR LIBERTYLINK® CANOLA

Tank Mix Partner	Rate (fl oz/A)
Assure® II	4 - 5 fl oz/A
Poast®	6 - 8 fl oz/A
Select® 2EC	2 – 3 fl oz/A
Select Max™	4 - 6 fl oz/A

APPLICATION RATE AND TIMING FOR CANOLA FOR TRANSGENIC SEED PROPAGATION

Up to three applications of Reckon 280SL at up to 22 fl oz/A per application may be made to canola for transgenic seed propagation. Applications may be made from the cotyledon stage up to the early

bolting stage (e.g., BBCH 18-30, between just prior to stem elongation/bolting, eight or more leaves and beginning of stem elongation, no internodes).

RESTRICTIONS TO THE DIRECTIONS FOR CANOLA FOR TRANSGENIC SEED PROPAGATION

- **DO NOT** apply than three applications of Reckon 280SL at up to 22 fl oz/A per application per year.
- **DO NOT** apply more than 66 fl oz/A of Reckon 280SL per year.
- **DO NOT** apply Reckon 280SL beyond the early bolting stage or within 65 days of harvesting canola seed.
- **DO NOT** use treated canola seed for food, feed or oil purposes.
- **DO NOT** apply Reckon 280SL if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.

APPLICATION DIRECTIONS FOR USE ON FIELD CORN AND SILAGE CORN

Apply Reckon 280SL only to corn labeled as LibertyLink[®]. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION RATE AND TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimal yield, early season weed removal is important.

Applications of Reckon 280SL on corn may be made with over-the-top broadcast or drop nozzles from emergence until corn is 24 inches tall or in the V-7 stage of growth, i.e., 7 developed collars, whichever comes first. For corn 24 inches to 36 inches tall, only apply Reckon 280SL using ground application and drop nozzles and avoid spraying into the whorl or leaf axils of the corn stalks. Applications of Reckon 280SL following the use of soil-applied insecticides will not injure corn.

Apply Reckon 280SL at 22 fl oz/A per application. A second application of Reckon 280SL or a tank mix application with a residual herbicide will be needed to control weeds that have not yet emerged at the time of application.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON FIELD CORN, AND SILAGE CORN

- **DO NOT** apply Reckon 280SL within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.
- **DO NOT** apply more than two applications of Reckon 280SL to the corn crop. Sequential applications should be at least 10 days apart.
- **DO NOT** apply more than 44 fl oz/A of Reckon 280SL on corn per year. If Reckon 280SL was used in a burndown application, no post emergence applications may be applied to the crop.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- **DO NOT** apply Reckon 280SL if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the "**Rotational Crop Restrictions**" section under the "**Information**" heading of this label for the appropriate rotational crop plant back intervals.

SPRAY ADDITIVES

For corn, Reckon 280SL must be applied with ammonium sulfate (AMS). It is recommended to use only fine feed grade or spray grade AMS at 3 lbs per acre (17 lbs/100 gallons). When temperatures exceed 85 °F, the rate of AMS can be reduced to 1.5 lbs per acre (8.5 lbs/100 gallons) to reduce potential leaf burn. Use of additional surfactants or crop oils may increase risk of crop response.

CORN TANK MIX INSTRUCTIONS

Certain herbicide tank mixes may aid in the performance of Reckon 280SL. No additional surfactant is needed with any tank mix partner. Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the corn to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing.

TANKMIX PARTNERS FOR RECKON 280SL ON LIBERTY LINK® CORN:

2,4-D	Halex GT	Pendimethalin ¹
acetochlor	Hornet® WDG	Permit®
Aim™ ²	Impact®	Python®WDG
Atrazine	Laudis®	s-metolachlor ²
Callisto™	Lexar® ²	Spirit®
Camix® ²	Lumax® ²	Status®
Capreno®	Metolachlor ²	Yukon®
Distinct®	nicosulfuron	Zemax
Guardzman Max®	NorthStar™	

¹Tankmixing with pendimethalin may result in reduced control of barnyardgrass, fall panicum, field sandbur, yellow foxtail, and volunteer corn.

²It is recommended that these products are tankmixed at half the use rate with Reckon 280SL to reduce risk of crop response. It is recommended that these products are tank mixed at ½ the use rate with Reckon 280SL to reduce risk of crop response.

CORN INSECTICIDE TANK MIX PARTNERS FOR RECKON 280SL:

To provide weed and insect control in corn, Reckon 280SL may be mixed with the following insecticides:

Ambush® Insecticide	Tombstone™ Helios®	Pounce® 3.2EC Insecticide
Asana® XL Insecticide	Lorsban® 4E Insecticide	Warrior™ Insecticide
Baythroid® XL Insecticide	Tombstone™	

APPLICATION DIRECTIONS FOR USE ON COTTON

Uniform, thorough spray coverage is necessary to achieve consistent weed control. Reckon 280SL may be applied as a broadcast, over-the-top, post-emergence spray or as a directed spray only to LibertyLink® cotton. This product may be applied post-emergence to non-LibertyLink® cotton varieties or cultivars by using equipment designed to minimize contact of the spray with the cotton foliage. See the Application Methods on Non- LibertyLink® Cotton section for selection of shielding equipment. Severe injury or death may result if the Reckon 280SL contacts the foliage or stems of cotton NOT labeled as LibertyLink®.

APPLICATION RATE AND TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimal yield, early season weed removal is important.

Apply Reckon 280SL to cotton from emergence up to the early bloom stage at 22 to 29 fl oz/A. Should environmental conditions prevent a timely herbicide application, a single application of up to 43 fl oz/A of Reckon 280SL may be made to cotton. If more than 29 fl oz/A are used in any single application, the yearly total may not exceed 72 fl oz/A, including all application timings. See Restrictions to the Directions for use on Cotton below for additional information.

Refer to the Weed Control Table for Row Crops section of this label for selection of the proper rate dependent upon weed species present and size. In weed populations with mixed species, select the highest rate required to control all the species. Volunteer LibertyLink® crop plants (corn, cotton, soybean, sugar beet) from the previous season will not be controlled by applications of Reckon 280SL.

A repeat application of Reckon 280SL or tank mixes with a residual herbicide will be needed to control weeds that have not emerged at the time of application. See the Tank Mix Instructions for Use on Cotton to select suitable tank mix partners.

Use Pattern	1st Application	2nd Application	3rd Application	Yearly Maximum
Option 1	22 - 29 fl oz/A	22 - 29 fl oz/A	22 - 29 fl oz/A	87 fl oz/A
Option 2	30 - 43 fl oz/ A	22 - 29 fl oz/A	None	72 fl oz/A

RESTRICTIONS TO THE DIRECTIONS FOR USE ON COTTON

- **DO NOT** apply Reckon 280SL to cotton in Florida, South of Tampa (Florida Route 60), or in Hawaii, except for test plots or breeding nurseries.
- **DO NOT** apply Reckon 280SL within 70 days prior to cotton harvest.
- Up to three applications of Reckon 280SL may be made to cotton per year at a maximum application rate of 29 fl oz/A. **DO NOT** apply more than 87 fl oz (including all application timings) to cotton per year under this application scenario. Sequential applications should be at least 10 days apart.

- If environmental conditions prevent timely applications resulting in large weeds or heavy infestations, a single application of Reckon 280SL at up to 43 fl oz/A may be made to cotton. **DO NOT** apply more than 43 fl oz of Reckon 280SL in a single application under this use scenario. If a single application greater than 29 fl oz is made, a subsequent application not to exceed 29 fl oz may be made to cotton. The yearly total use rate under this scenario may not exceed 72 fl oz of Reckon 280SL. Sequential applications should be at least 10 days apart.
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the "**Rotational Crop Restrictions**" section under the "**Information**" heading of this label for the appropriate rotational crop plant back intervals.

APPLICATION METHODS TO LIBERTYLINK® COTTON

Refer to the Weed Control Table for Row Crops to select the proper application rate based upon the weeds present and their size. Uniform and thorough spray coverage is required to achieve consistent weed control. For ground application, apply Reckon 280SL to LibertyLink® cotton as an over-the-top foliar spray or as a spray directed to the lower one-third of the cotton stand.

APPLICATION METHODS TO NON-LIBERTYLINK COTTON

Application of Reckon 280SL to cotton varieties not labeled as LibertyLink® requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. A hooded sprayer directs the spray onto weeds, while shielding the cotton stand from contact. Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid exposure of the desirable vegetation to the spray.

With a hooded sprayer, the spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. If the hoods are raised, spray particles may escape and come into contact with the cotton, causing damage or destruction of the crop.

Herbicide rates and spray volume instructions are presented as broadcast equivalents and must be reduced in proportion to the area actually treated. Use the following formulas to calculate the correct rate and volume per planted (field) acre:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Amount of banded product needed per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast spray VOLUME per acre} = \text{Banded Spray Volume needed per acre}$$

POST-HARVEST

Reckon 280SL may be applied as a post-harvest burndown treatment to fields (after cotton harvest). Up to 43 fl oz/A of Reckon 280SL may be applied in a single application to control larger weeds growing in the crop at the time of harvest. If more than 29 fl oz/A is used in a single application, the yearly total may not exceed 72 fl oz/ A, including all application timings. Refer to the Rotational Crop Restrictions section of this label for appropriate rotational crop information.

COTTON TANK MIX INSTRUCTIONS

Certain tank mixes may aid in the performance of Reckon 280SL. No additional surfactant is needed with any tank mix partner. Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing.

LibertyLink® Cotton: For cotton tolerant to Reckon 280SL, Dual Magnum® or Staple® Herbicide may be tank-mixed with Reckon 280SL and applied over-the-top post-emergence to enhance weed control and/or provide residual control.

All Cotton Types: The following herbicides may be mixed with Reckon 280SL for hooded-spray application to enhance weed control and/or provide residual weed control:

Aim™	Cotoran® DF	Dual Magnum®	Pendimax™ 3.3	Staple®
Caparol® 4L	Direx® 4L	Glyphosate	Prowl® 3.3EC	
Cotoran® 4L	Direx® 80DF	Karmex® DF	Select Max™	

POSTEMERGENCE OVER-THE-TOP TANKMIX PARTNERS FOR RECKON 280SL ON LIBERTYLINK® COTTON

Assure II	metolachlor	Clethodim
Poast Plus	Fusilade DX	Select Max
Fusion	Staple	

APPLICATION DIRECTIONS FOR USE ON SOYBEANS

Apply Reckon 280SL only to soybean designated as LibertyLink®. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

APPLICATION RATE AND TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Adding ammonium sulfate with Reckon 280SL may improve weed control if weeds are under stress. For optimal yield, early season weed removal is important.

Applications of Reckon 280SL on soybeans may be made from emergence up to but not including the bloom growth stage.

Apply Reckon 280SL to LibertyLink® soybeans from emergence up to but not including the bloom growth stage at 22 to 29 fl oz/A. See weed chart to determine rate. Should environmental conditions prevent a timely herbicide application, a single application of up to 36 fl oz/A of Reckon 280SL may be made to soybeans followed by one additional application at a maximum of 29 fl oz/A with a yearly maximum of 65 fl oz/A. Reckon 280SL may be applied alone, or in a tank mix application with a residual herbicide to control weeds that have not yet emerged at the time of application.

Although timely post applications of Reckon 280SL can provide complete weed control, residual herbicides at burndown, planting, or tank mixed with Reckon 280SL help ensure optimal weed management, particularly if environmental conditions delay timely post applications. Residual herbicides can also reduce early season weed competition and are a key element of good weed resistance management practices.

Use Pattern Rate Ranges		
1 st Application	2 nd Application	Yearly Maximum
22 – 36 fl oz/A	22 – 29 fl oz/A	65 fl oz/A

RESTRICTIONS TO THE DIRECTIONS FOR USE ON SOYBEANS

- **DO NOT** apply Reckon 280SL within 70 days of harvesting soybean seed.
- **DO NOT** apply more than 65 fl oz/A of Reckon 280SL on soybeans per year.
- **DO NOT** apply more than 36 fl oz/A of Reckon 280SL in a single application.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- **DO NOT** apply Reckon 280SL if soybeans show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the "**Rotational Crop Restrictions**" section under the "**Information**" heading of this label for the appropriate rotational crop plant back intervals.
- Sequential applications should be at least 5 days apart.

SOYBEAN TANK MIX INSTRUCTIONS

Certain herbicide tank mixes may complement Reckon 280SL. No additional surfactant is needed with any tank mix partner. Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the soybean to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing.

TANKMIX PARTNERS FOR RECKON 280SL IN LIBERTYLINK® SOYBEANS

Assure® II	Fusion®	Raptor™
Classic®	Harmony®	Reflex®
clethodim	Optill	Resource®
Cobra®	metolachlor	Select Max®
Fierce	Phoenix™	Sharpen
First Rate®	Poast Plus®	Synchrony®XP
Flexstar®	Prefix	Ultra Blazer®
Fusilade® DX	Pursuit®	

APPLICATION DIRECTIONS FOR CANOLA, CORN, COTTON, AND SOYBEAN SEED PROPAGATION

Reckon 280SL may be applied to select out susceptible "segregates," i.e., canola, corn, cotton, and soybean plants that are not tolerant to glufosinate-ammonium during seed propagation.

- Canola:** Reckon 280SL may also be used in canola seed propagation as a foliar spray to selectively eliminate canola plants that do not carry a gene that imparts tolerance to glufosinate-ammonium and as such, can be applied to remove susceptible segregates during canola seed propagation. Breeding material not possessing the glufosinate-ammonium tolerance gene will be severely injured or killed if treated with this herbicide. See **Application Use Directions for Use on Canola** for use rates and application timing.
- Corn:** Inbred lines, plants not possessing glufosinate-ammonium tolerance, will be severely injured or killed if treated with this herbicide. A hooded sprayer may be used to protect plants from coming into contact with the herbicide application. For the selection of tolerant corn "segregates", Reckon 280SL may be applied at 22 fl oz/A plus AMS at 3 lb/A (17 lb/100 gallons) when corn is in the V-3 to V-4 stage of growth, i.e., 3 to 4 developed collars. A second treatment of 22 fl oz/A plus AMS at 3 lb/A may be applied when the corn is in the V-6 to V-7 stage of growth or up to 24" tall. Sequential applications should be at least 10 days apart. When temperatures exceed 85°F, the rate of AMS can be reduced to 1.5 lbs/A (8.5 lb/100 gallons) to reduce potential leaf burn.
- Cotton:** Reckon 280SL may also be used in cotton seed propagation as a foliar spray to selectively eliminate cotton plants that do not carry a gene that imparts tolerance to glufosinate-ammonium and as such, can be applied to remove susceptible segregates during cotton seed propagation. Breeding material not possessing the glufosinate-ammonium tolerance gene will be severely injured or killed if treated with this herbicide. See **Application Use Directions for Use on Cotton** for use rates and application timing.
- Soybean:** For the selection of tolerant soybean "segregates", Reckon 280SL may be applied at up to 22 to 36 fl oz/A when soybean is in the third trifoliolate stage. A second treatment of 22 to 29 fl oz/A may be applied up to but not including the bloom growth stage of soybean. Sequential applications should be at least 5 days apart.

Grass Weeds Controlled by Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Height)		Comments on Weed Growth Stages/Application Timing/Number of Applications
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)	
Barley, volunteer	1 – 2 leaf (2")	3 leaf (3")	Multiple applications may be required
Barnyardgrass	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Corn, volunteer	1 – 2 leaf (2")	3 – 4 leaf (6")	---
Crabgrass, large	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Crabgrass, smooth	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Cupgrass, woolly	1 – 5 leaf (4")	(8")	---
Foxtail, giant	1 – 4 leaf (4")	5 – 6 leaf (4")	Maximum of 2 tillers
Foxtail, green	1 – 4 leaf (4")	5 – 6 leaf (4")	Maximum of 2 tillers
Foxtail, yellow	1 – 3 leaf (2")	4 leaf (2")	Apply prior to tillering
Millet, volunteer proso	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Millet, wild proso	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Oat, wild	1 – 2 leaf (2")		Maximum of 1 tiller
Panicum, fall	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Panicum, Texas	1 – 3 leaf (2")	4 – 5 leaf (3")	Maximum of 1 tiller
Sandbur, field	---	1 – 4 leaf (2")	Apply prior to tillering
Wheat, volunteer	1 – 2 leaf (2")	3 leaf (3")	Maximum of 1 tiller

***Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.**

For improved control of heavy populations or larger than recommended volunteer wheat, volunteer barley, yellow foxtail and wild oats, Reckon 280SL can be tank mixed with Assure® II Herbicide, Poast® Herbicide, Prism® Herbicide or Select® 2EC Herbicide.

Perennial Weeds Controlled by Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Height/Diameter)		Comments on Weed Growth Stages/Application Timing/Number of Applications
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)	
Quackgrass	---	1 – 3 leaf (3")	Multiple applications required
Sowthistle, perennial	---	1 – 4 leaf (3")	Multiple applications required
Thistle, Canada	---	1 – 4 leaf (3")	Multiple applications required

***Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.**

Broadleaf Weeds Controlled by Reckon 280SL

Weed Species	Growth Stage of Weed* (Maximum Diameter)	
	15 fl oz/A (0.9 pt/A)	20 fl oz/A (1.25 pt/A)
Buckwheat, wild	1 – 4 leaf (2")	5 – 6 leaf (3")
Buffalobur	1 – 4 leaf (2")	5 – 6 leaf (3")
Carpetweed	---	1 – 4 leaf (2")
Chickweed, common	1 – 4 leaf (2")	5 – 6 leaf (3")
Cocklebur, common	1 – 6 leaf (3")	7 – 8 leaf (5")
Kochia	(1")	(2")
Ladysthumb	1 – 2 leaf (1")	3 – 4 leaf (3")
Lambsquarter, common	1 – 2 leaf (1")	4 – 5 leaf (3")
Mallow, Venice	1 – 4 leaf (2")	5 – 6 leaf (3")
Marshelder	1 – 2 leaf (1")	3 – 4 leaf (2")
Mustard, wild	1 – 4 leaf (2")	5 – 6 leaf (3")
Nightshade, eastern black	1 – 4 leaf (2")	5 – 6 leaf (3")
Pigweed, prostrate	(1")	(3")
Pigweed, redroot	1 – 2 leaf (1")	3 – 4 leaf (3")
Pigweed, smooth	1 – 2 leaf (1")	3 – 4 leaf (3")
Pigweed, spiny	1 – 2 leaf (1")	3 – 4 leaf (3")
Purslane, common	(1")	(2")
Ragweed, giant	1 – 4 leaf (2")	5 – 6 leaf (3")
Shepherd's purse	1 – 4 leaf (2")	5 – 6 leaf (3")
Smartweed, Pennsylvania	1 – 2 leaf (1")	3 – 4 leaf (3")
Sowthistle, annual	1 – 4 leaf (2")	5 – 6 leaf (3")
Sunflower, common	1 – 6 leaf (3")	7 – 8 leaf (5")
Thistle, Russian	(1")	(2")
Velvetleaf	1 – 2 leaf (1")	3 – 4 leaf (3")

*Apply up to 30 fl oz/A (1.88 pt/A) if weeds exceed the growth stage shown in the table.

APPLICATION DIRECTIONS FOR USE ON SWEET CORN*

Apply Reckon 280SL only to corn labeled as LibertyLink®.

*Not for use in CA

Application Timing for Sweet Corn:

Applications of Reckon 280SL on sweet corn may be made from emergence until sweet corn is 24" tall or the V-7 stage of growth; i.e., 7 developed collars, whichever comes first. Apply at a rate of 20 fl oz/A. Reckon 280SL must be applied with ammonium sulfate (AMS) for use on sweet corn. Two applications of Reckon 280SL can be made to sweet corn per year.

Restrictions to the Directions for Use on Sweet Corn:

- **DO NOT** apply Reckon 280SL within 50 days of harvesting sweet corn ears and within 55 days of harvesting stover.
- **DO NOT** apply more than 40 fl oz/A of Reckon 280SL on sweet corn per year.
- **DO NOT** apply more than two applications of Reckon 280SL to the sweet corn crop. Sequential applications should be at least 10 days apart.

- If Reckon 280SL was used in a burndown application, no postemergence applications may be made to the crop.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- **DO NOT** apply Reckon 280SL if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall etc.).
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the **Rotational Crop Restrictions** section under the **Product Information** heading of this label for the appropriate rotational crop plantback intervals.

See the **Application Methods for Broadcast Application, Mixing Instructions and Weed Control for Row Crops** tables on this label for further instruction.

Tank Mix Instructions for Use on Sweet Corn:

Reckon 280SL may be tankmixed with Laudis® Herbicide, Callisto® Herbicide, Atrazine or Permit®. When using Reckon 280SL in tankmix combinations, carefully follow the Directions for Use labeling of the selected partner.

APPLICATION DIRECTIONS FOR USE ON LISTED TREE, VINE, AND BERRY CROPS

Apply Reckon 280SL to the tree, vine, and berry crops listed below. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

REGISTERED CROPS

- Bushberries: blueberry, currant, elderberry, gooseberry and huckleberry
- Other Berries: Lingonberry, juneberry and salal
- Citrus: lemon, orange, grapefruit, lime, mandarin, tangerine, tangelo, calamondin, kumquat, pummelo, citron and tangor; cultivars, varieties and/or hybrids of these
- Olives
- Pome Fruits: apples, pears, crabapple, loquat, mayhaw, quince, azarole, medlar and tejocote; cultivars, varieties and/or hybrids of these
- Stone Fruits: apricot, cherry, peach, nectarine, plum, capulin, jujube and sloe; cultivars, varieties and/or hybrids of these
- Tree Nuts: almonds, filberts, hickory nuts, macadamia nuts (bush nuts), pecans, pistachios, and walnuts
- Vineyards: all grape varieties (table, wine, and raisins)

APPLICATION RATE AND TIMING

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Reckon 280SL. Refer to the Weed Control for Row Crops section of this label for selection of the proper rate dependent upon weed species present and size. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. Do not retreat these weeds with Reckon 280SL until sufficient regrowth has occurred.

Apply Reckon 280SL as a directed spray to control undesirable vegetation in tree, vine and berries listed on this label. Apply as a broadcast, banded, or spot treatment application depending on the

situation to control weeds listed under the heading “Weeds Controlled in Tree, Vine and Berry crops”. Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of Reckon 280SL may be necessary to control plants generating from underground parts or seed.

Avoid contact of Reckon 280SL solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees and vines. **Only trunks with callused, mature brown bark should be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of Reckon 280SL with parts of trees or vines other than mature brown bark can result in serious damage.**

Application Methods for Broadcast Applications

Apply Reckon 280SL at the rates listed below for broadcast applications based on weed size and stage of growth.

Weed Size and Stage	Reckon 280SL Rate
Weeds < 3” in height	48 fl oz/A
Weeds < 6” in height pre-tiller grasses	56 fl oz/A
Weeds > 6” in height and/or grasses that have tillered	56-82 fl oz/A

Application Methods for Banded Spray Applications

Banded applications may be used using the following formula to calculate the amount of herbicide needed for orchard or vineyard strip sprays:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre broadcast} = \text{Amount of herbicide needed for treatment}$$

Application Methods for Spot or Directed-Spray Applications

For spot or directed spray application, mix Reckon 280SL at 1.7 fl oz of product per gallon of water. Apply to undesirable vegetation foliage until wet but prior to runoff. Ensure uniform and complete coverage. Thoroughly clean the sprayer following use. DO NOT make spot or directed spray applications to tree or vine trunk as injury may occur.

Weeds Controlled in Tree, Vine and Berry Crops

Broadleaf Weeds

Alkali sida	Fleabane, annual	Morningglory, entireleaf	Redmaids
Ammannia, purple	Goosefoot	Morningglory, ivyleaf	Shepherd's-Purse
Arrowhead, California	Gromwell, field	Morningglory, pitted	Smartweed, Pennsylvania
Buckwheat, wild	Groundcherry, cutleaf	Mullein, turkey	Sowthistle, annual
Buffalobur	Groundsel, common	Mustard, wild	Spurge, prostrate
Burclover, California	Henbit	Nettle	Starthistle, yellow
Carpetweed	Jimsonweed	Nightshade, black	Sunflower, common
Chickweed, common	Knotweed	Nightshade, eastern	Sunflower, prairie
Chinese thornapple	Kochia	Nightshade, hairy	Sunflower, volunteer
Cocklebur, common	Lambsquarters, common	Pennycress	Swinecress
Cudweed	Lettuce, miner's	Pigweed, redroot	Thistle, Russian
Cutleaf	Lettuce, prickly	Pineapple-weed	Turnip, wild
eveningprimrose	London rocket	Puncturevine	Velvetleaf
Dodder	Mallow, common	Purslane, common	Vervain
Eclipta	Malva (little mallow)	Radish, wild	Vetch
Fiddleneck	Marestail	Ragweed, common	Virginia copperleaf
Filaree	Mayweed	Ragweed, giant	Willowherb, panicle
Filaree, redstem			

Grass Weeds

Barnyardgrass	Crabgrass, smooth	Junglerice	Shattercane
Bluegrass, annual	Cupgrass, woolly	Oat, wild	Sprangletop
Brome, ripgut	Foxtail, giant	Panicum, fall	Stinkgrass
Bromegrass, downy	Foxtail, green	Panicum, Texas	Wheat, volunteer
Canarygrass	Foxtail, yellow	Rush, toad**	Windgrass
Chess, soft	Goosegrass	Ryegrass, annual	Witchgrass
Crabgrass, large	Johnsongrass, seedling	Sandbur, field	

Biennial and Perennial Weeds

Aster, white heath	Dallisgrass	Mullein, common	Rocket, yellow
Bindweed, field	Dandelion	Mustard, tansy	Rose, wild
Bindweed, hedge	Dock, curly	Nutsedge, purple	<i>Rubus</i> spp.
Bluegrass, Kentucky	Dogbank (hemp)	Nutsedge, yellow	Spurge, leafy
Bromegrass, smooth	Fescue	Onion, wild	Thistle, bull
Bulrush**	Goldenrod, gray	Orchardgrass	Thistle, musk
Burdock	Guineagrass	Paragrass	Torpedograss
Canada thistle	Horsetail	Plantain	Vaseygrass
Clover, Alsike	Lovegrass	Poison ivy/oak	Woodsorrel
Clover, red	Mugwort	Quackgrass	Yarrow, common
Clover, white			

**indicates suppression

RESTRICTIONS TO THE DIRECTIONS FOR USE ON TREE, VINE, AND BERRY CROPS

1. **DO NOT** apply more than 164 fl oz of Reckon 280SL per acre (3 lbs ai/A) to berry bushes in a 12-month period.
2. **DO NOT** apply more than 246 fl oz of this product per acre to tree nuts, vines, and tree fruits in any calendar year.
3. **DO NOT** graze, harvest, and/or feed treated orchard cover crops to livestock.
4. **DO NOT** apply this product through any type of irrigation system.
5. **DO NOT** apply this product aerially to tree, berry, or vine crops.
6. **DO NOT** apply this product within 14 days of nut, apple, berry or grape harvest.
7. **DO NOT** make spot spray applications to apple suckers, as tree injury may occur.

SUCKER CONTROL WITH RECKON 280SL

Reckon 280SL will reduce or eliminate sucker growth when applied to suckers that are young, green and uncallused. For sucker control, apply a split application approximately 4 weeks apart at 56 fl oz/A. Coverage of all sucker foliage is necessary for optimum control. Suckers should not exceed 12 inches in length.

TANK MIX PARTNER

Reckon 280SL does not provide residual weed control or control of unexposed plant parts. Certain herbicide tank mixes may aid in the performance of Reckon 280SL or be added to provide residual herbicide activity. No additional surfactant is needed with any tank mix partner. Reckon 280SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Reckon 280SL cannot be mixed with any product containing a label prohibition against such mixing.

Chateau	Karmex® DF	Simazine 80W	Solicam® DF
Devrinol® 50WP	Princep® 4L	Simazine 90	Surflan® A.S.
Goal® 1.6E	Simazine 4L	Sinbar® 80W	

APPLICATION DIRECTIONS FOR POTATO VINE DESICCATION

APPLICATION RATE AND TIMING

Apply Reckon 280SL at the beginning of natural senescence of potato vines. Apply 21 fl oz/A. Do not split this application or apply more than one application per harvest. Potato varieties with heavy or dense vines may require an application of another desiccation product to complete vine desiccation.

Thorough coverage of the potato vines to be desiccated is essential. Use a sufficient volume of water (20 to 100 gpa) to obtain a thorough coverage of the potato vines. Vary the gallons of water per acre and the spray pressure as indicated by the density of the potato vines to assure thorough spray coverage. Increase the spray volume to at least 30 gallons of water per acre when the potato vine canopy is dense or under cool and dry conditions. Apply Reckon 280SL with the spray boom as low as possible to achieve thorough coverage of the potato vines for best control and to minimize drift potential.

RESTRICTIONS TO THE DIRECTIONS FOR USE IN POTATO VINE DESICCATION

1. **DO NOT** apply more than 21 fl oz/A to potato vines per year.
2. **DO NOT** harvest potatoes until 9 days or more after application of Reckon 280SL.
3. **DO NOT** apply to potatoes grown for seed.
4. Sweet corn and cotton may be planted at any time after the application of Reckon 280SL as a potato vine desiccant.
5. **DO NOT** plant treated areas to wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale until 30 or more days after an application of Reckon 280SL as a potato vine desiccant.
6. **DO NOT** plant treated areas to crops other than those listed in this use precautions section until 120 or more days after an application of Reckon 280SL as a potato vine desiccant.

APPLICATION DIRECTIONS FOR USE ON RICE*
--

*Not for use in CA

THOROUGH SPRAY COVERAGE IS VERY IMPORTANT. For best results, apply to emerged, young, actively growing weeds. Reckon 280SL is a foliar active material with little or no soil residual activity. Weeds that emerge after application will not be controlled. Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present, or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness. Reckon 280SL is rainfast 4 hours after application to most weed species. Rainfall within 4 hours after application may necessitate retreatment or reduced weed control may result.

RESTRICTIONS TO THE DIRECTIONS FOR USE ON RICE

- **DO NOT** exceed 48 fl oz of Reckon 280SL per year.
- **DO NOT** apply Reckon 280SL within 70 days of harvesting rice.
- **DO NOT** plant rotation crops in a field treated with Reckon 280SL within 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product. The crops listed on this label may be planted at any time.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** use paddy water from a rice field treated with Reckon 280SL for irrigation, or as a water source for livestock or for raising crayfish.
- **DO NOT** add surfactants or crop oils. A silicon-based anti-foam agent may be added if needed.

APPLICATION TIMING FOR THE SOUTHERN UNITED STATES (ARKANSAS, LOUISIANA, MISSISSIPPI, MISSOURI, TEXAS)

Applications of Reckon 280SL on rice may be made from the 1-leaf stage through the mid-tillering stage of development. Refer to the **Rate Tables for Weed Control in Rice** to select the proper rate to

use to control the weed species present. Reckon 280SL will have an effect on weeds that are larger than the recommended leaf stage, however, speed of activity and control may be reduced.

Rice fields should be as level as possible and free of large clods to obtain uniform germination of rice and grassy weeds and to ensure uniform flood levels. If necessary, fields may be flushed prior to treatment so that the rice and grass/broad leaf weeds are actively growing at the time of treatment. If the rice field is flushed, allow sufficient time for germination of the weed species to occur prior to treatment.

Apply Reckon 280SL prior to the permanent flood when weeds are in the 1- to 5-leaf stage. A second application is recommended after a new flush of weeds emerge. A second application may be made from 10 -14 days after the first application up to the mid-tillering growth stage of the rice. For optimum weed control, apply Reckon 280SL before canopy closure to ensure thorough spray coverage of the weed species.

When applying Reckon 280SL post flood, lower the water level so that 75 of the weed foliage is exposed. The water level may be brought back to normal 48 hours after the herbicide application.

Rate Tables for Weed Control in Rice

1. *Southern United States (Arkansas, Louisiana, Mississippi, Missouri, Texas)*

Grass Weeds Controlled with Reckon 280SL in Rice Grown in the Southern United States

Weed Species	Maximum Weed Growth Stage (leaf/tiller)	
	20 fl oz/A	24 fl oz/A
Barnyardgrass	4 leaf	2 tiller
Crabgrass, large	4 leaf	2 tiller
Fall Panicum	4 leaf	2 tiller
Johnsongrass	4 leaf	2 tiller
Rice, red*	4 leaf	2 tiller
Signalgrass, broadleaf	4 leaf	2 tiller
Sprangletop	4 leaf	2 tiller
Watergrass	6 leaf	2 tiller

*For optimum red rice control, make two applications of Reckon 280SL. The first application should be made when the red rice is in the 2 – 3 leaf stage. The second application should be made after the newly emerged red rice reaches the 2 – 3 leaf stage but before the white rice reaches the mid-tillering stage of development.

Broadleaf Weeds Suppressed or Controlled with Reckon 280SL In Rice Grown in the Southern United States

Weed Species	Maximum Weed Height in Diameter (inches)	
	20 fl oz/A	24 fl oz/A
Ammania	2	4
California Arrowhead	**	4
Cocklebur, common		10
Curly Indigo	6	8

Dayflower	2	4
Eclipta	2	6
Morningglory, ivyleaf	4	8
Morningglory, pitted	4	8
Northern jointvetch	4	8
Pennsylvania Smartweed	4	8
Sesbania hemp	4	10

**indicates suppression

Reckon 280SL applied at 24 fl oz/A may control or suppress the sedges shown in the following table. Control of sedges may be enhanced by using a second application or by a tank mix with other herbicides recommended on this label.

Sedges suppressed with Reckon 280SL in Rice Grown in the Southern United States

Sedges	24 fl oz/A
Bulrushes	**
Flatsedge	**
Nutsedge	**
Smallflower Umbrellaplant	**

**indicates suppression

TANK MIX INSTRUCTIONS FOR USE IN RICE

When using Reckon 280SL in tank mix combinations, follow the precautions and directions of the most restrictive label for the appropriate timing, rate and crop response information.

1. Southern United States (Arkansas, Louisiana, Mississippi, Missouri, Texas)

To enhance weed control and/or provide residual control in rice, Reckon 280SL may be mixed with the following herbicides:

Arrosolo® 3.3E Herbicide

Basagran® Herbicide

Bolero® EC Herbicide

Londax® Herbicide

Prowl® 3.3EC Herbicide

Propanil

Stam® Herbicide

Permit® Herbicide

APPLICATION DIRECTIONS FOR USE IN RICE SEED PROPAGATION*

*Not for use in CA

Reckon 280SL is to be applied as a foliar spray to selectively remove susceptible segregates (i.e., undesirable rice plants which are not tolerant to glufosinate-ammonium and to control a broad spectrum of emerged grass and broadleaf weeds in rice seed production fields). Inbred lines or breeding material not possessing the glufosinate-ammonium tolerance gene will be severely injured or killed if treated with this herbicide. Apply Reckon 280SL exclusively to rice seed propagation fields in which the desired plants are glufosinate-ammonium tolerant.

THOROUGH SPRAY COVERAGE IS VERY IMPORTANT. Reckon 280SL works best when weeds are small and the crops and weeds are actively growing. Visual effects and control of rice susceptible segregates from Reckon 280SL applications occur within 2 to 4 days after application under good growing conditions. The ability of Reckon 280SL to eliminate rice plants not tolerant to Reckon 280SL may be reduced when heavy dew, fog, or mist/rain is present on the crop, or when crop is under stress due to drought, cool temperatures, or extended periods of cloudiness.

Rice fields should be as level as possible and free of large clods to obtain uniform germination on rice and grassy weeds and to ensure uniform flood levels. If necessary, fields may be flushed prior to treatment. If fields are flushed prior to treatment, flush in sufficient time so that the rice and grass/broadleaf weeds are actively growing at time of treatment.

Do not allow spray to contact foliage or green tissue of desirable vegetation other than rice lines in which the desired plants are glufosinate-ammonium tolerant. This product will injure any other green vegetation contacted by the spray.

INSTRUCTIONS FOR SEED HANDLING, STORAGE AND USE

Seed from treated plants must be held in secured storage until used for breeding of glufosinate-ammonium tolerant rice seed, or destroyed. Seed from treated plants must be labeled as follows: Do Not Use for Feed or Food Purposes. Store Away from Feed and Foodstuffs. In addition, label the seed with the "Seed Disposal" statements found in the "Storage and Disposal" section of this label.

RESTRICTIONS TO THE DIRECTIONS FOR USE

1. **DO NOT** use rice, any rice possessed commodities or rice straw treated with Reckon 280SL for food or feed consumption.
2. **DO NOT** exceed 80 fl oz/A of Reckon 280SL per year on rice being treated for segregate control in seed production fields.
3. **DO NOT** plant rotation crops in a field treated with Reckon 280SL for 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product.
4. **DO NOT** apply this product through any type of irrigation system.

Rate instructions and Timing for Seed Production

For the selection of susceptible rice (segregates) Reckon 280SL must be applied at 40 fl oz/A when rice is in the 1 to 3 leaf stage of growth. A second treatment of 40 fl oz/A must be applied 10 days later or up until the rice is in the mid-tillering state of growth.

- **DO NOT** exceed 80 fl oz (1.46 lbs ai/A) per single application.
- 2 applications can be made at 40 fl oz (0.73 lb ai/A) with a minimum 10 day re-treatment interval.
- **DO NOT** exceed 80 fl oz (1.46 lbs ai/A) per year.
- Minimum paddy depth of 4 inches.
- If 1 application of 80 fl oz is made, the application must be made to a dry field. A minimum 7 day holding period after flooding of the field is required.
- If 2 applications are made, the first application must be made to a dry field.
- The second application may be made to a flooded field with a required 55 day holding period for a 4 inch paddy depth or a 30 day holding period for an 8 inch paddy depth.

WATER MANAGEMENT

A sufficient portion of the target grassy weed plant must be exposed to Reckon 280SL for satisfactory control to be achieved. Therefore, if necessary, lower or allow water to recede so that at least 75 of the weed foliage is exposed above the water level. Do not increase the water level for at least 48 hours following the application of Reckon 280SL. The water level may be brought back to normal level following this period.

TANK MIX INSTRUCTIONS FOR RECKON 280SL USE IN RICE SEED PROPAGATION

When using Reckon 280SL in tank mix combinations, follow the precautions and directions of the most restrictive label for the appropriate timing, rate, and crop response information.

1. Southern United States (Arkansas, Louisiana, Mississippi, Missouri, Texas)

To enhance weed control and/or provide residual control in rice, Reckon 280SL may be mixed with the following herbicides:

Arrosolo® 3 3E Herbicide

Basagran® Herbicide

Bolero® 8EC Herbicide

Londax® Herbicide

Prowl® 3.3EC Herbicide

Stam® Herbicide

Permit® Herbicide

FALLOW FIELDS OR POST HARVEST

Reckon 280SL may be used as a substitute for tillage in fallow fields to control or suppress weeds listed in the **Weed Control for Row Crops** section of this label. Applications may be made in fallow fields, post-harvest, prior to planting or emergence of any crop listed on this label. Apply Reckon 280SL at 22 or 29 fl oz/A to fallow fields to control specific weeds. Reckon 280SL must be applied with ammonium sulfate. Tank mixes with 2,4-D, glyphosate or atrazine are recommended with Reckon 280SL to enhance total weed control. When using Reckon 280SL in tank mix combinations, follow the precautions and directions of use of the most restrictive label. See the **Application and Mixing Procedures** section of this label for additional information on how to apply this product. See the "Information" section of this label for rotational crop restrictions.

FARMSTEADS, RECREATIONAL AND PUBLIC AREAS

When applied as listed, Reckon 280SL controls undesirable plant vegetation in non-crop areas around farmstead building foundations, shelter belts, along fences, airports, commercial plants, storage and lumber yards, educational facilities, fence lines, ditch banks, dry ditches, roadsides, schools, parking lots, tank farms, pumping stations, parks, other public areas and general nonselective farmstead weed control. Refer to the **Application Directions for Use on Listed Tree, Vine, and Berry Crops** section of this label for appropriate application broadcast and spot spray application rates and lists of weeds controlled.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well-ventilated place. Storage temperature should not exceed 125° F. If storage temperature for bulk Reckon 280SL is below 32° F, the material should not be pumped until its temperature exceeds 32° F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)]

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[All refillable container types (containers with capacities greater than 50 lbs)]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. This is a sealed returnable container to be used only for Reckon 280SL. When this container is empty, it must not be opened, cleaned, or discarded. Empty containers must be returned to the original purchase location.

[Bottom discharge Intermediate Bulk Container (IBC) (containers with capacities greater than 50 lbs)]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Empty the remaining contents from the Intermediate Bulk Container (IBC) into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inch on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve. Contact your Ag retailer or Orion GFS for container return, disposal and recycling recommendations.

SEED DISPOSAL: To dispose of out-of date or otherwise unmarketable seed from plants which have been treated with Reckon 280SL, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

CONDITIONS OF SALE AND LIMITED WARRANTY

The Directions for Use are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Orion GFS, LLC or the SELLER. To the extent consistent with applicable law, all such risks shall be assumed by the buyer.

Orion GFS, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. ORION GFS, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ORION GFS, LLC AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

ORION GFS, LLC and the SELLER offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of ORION GFS, LLC.

- Liberty, LibertyLink, Rely, Laudis, and the LibertyLink design are registered trademarks of Bayer CropScience.
- Impact is a registered trademark of Amvac Chemical Company.
- Guardsman Max, Poast, Prowl, Pursuit, and Status are registered trademarks and Distinct and Raptor are trademarks of BASF Corporation.
- Firstrate, Surflan, Goal, and Hornet are registered trademarks and Pendimax is a trademark of Dow AgroSciences.
- Assure, Classic, Direx, Harmony, Karmex, Staple, Synchrony, Sinbar, and Vitron are registered trademarks of E. I. DuPont de Nemours Company.
- Aim is a trademark of FMC.
- Cotoran is a registered trademark of Makhteshim Agan North America.
- Permit and Yukon are registered trademarks of Monsanto.

- Camix, Caparol, Dual Magnum, Flexstar, Fusilade, Fusion, Lexar, Lumax, Reflex, Solicam, Princep, and Spirit are registered trademarks and Callisto and NorthStar, are trademarks of Syngenta Group Company.
- Cobra, Resource, Chateau, and Select are registered trademarks and Phoenix and Select Max are trademarks of Valent U.S.A. Company.
- Ultra Blazer is a registered trademark of United Phosphorus, Inc.