

# Tebu-Crop 3.6F

**ACTIVE INGREDIENT:**

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha(1,1-dimethylethyl)-  
1H-1,2,4-triazole-1-ethanol ..... 38.7%

**OTHER INGREDIENTS:** ..... 61.3%

**TOTAL:** ..... **100.0%**

Contains 3.6 pounds tebuconazole per gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See complete First Aid, Precautionary Statements, and Directions For Use in booklet.

**STOP-Read the label before use.**

**Manufactured For:**

**Sharda USA LLC**



7217 Lancaster Pike, Suite A  
Hockessin, Delaware 19707

EPA Reg. No. 83529-11

EPA Est. No. 70815-GA-001

**Net Contents: 2.5 Gallons**

## FIRST AID

|                                |  |
|--------------------------------|--|
| <b>If Swallowed:</b>           | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul> |
| <b>If On Skin or Clothing:</b> | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |
| <b>If In Eyes:</b>             | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |
| <b>If Inhaled:</b>             | <ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>                                      |

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically.

**Symptoms of Poisoning:** The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

### EMERGENCY NUMBERS

For 24-hour medical emergency assistance (human or animal) call **1-800-222-1222**. For chemical emergency assistance (spill, leak, fire, or accident) call CHEMTREC at **1-800-424-9300**.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, barrier laminate, or butyl rubber  $\geq$  14 mils or nitrile rubber  $\geq$  14 mils or neoprene rubber  $\geq$  14 mils or polyvinyl chloride  $\geq$  14 mils or viton  $\geq$  14 mils
- Shoes plus socks

**User Safety Requirements:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Groundwater Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, barrier laminate or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or polyvinyl chloride ≥ 14 mils or viton ≥ 14 mils
- Shoes plus socks

### **AGRICULTURAL USE DIRECTIONS**

#### **SHAKE WELL BEFORE USING**

**RESTRICTIONS: Aerial application is prohibited in New York State.**

**Chemigation: Do not apply this product through any type of irrigation system.**

### **SPRAY DRIFT MANAGEMENT**

Do not allow this product to drift.

#### **Foliar Spray Drift Management**

Avoiding spray drift from foliar applications is the responsibility of the applicator. Similar to aerial spray drift, the interaction of many equipment- and weather-related factors determine the potential for spray drift from foliar applications. To protect water resources, the applicator and the grower are responsible for considering all these factors when making decisions.

## **Aerial Spray Drift Management**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the [Aerial Drift Reduction Advisory Information](#).

### **AERIAL DRIFT REDUCTION ADVISORY**

This section is advisory in nature and does not supersede the mandatory label requirements.

#### **INFORMATION ON DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity**, and **Temperature Inversions**).

#### **CONTROLLING DROPLET SIZE**

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure:** Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use high flow rate nozzles instead of increasing pressure.

- **Number of Nozzles:** Use minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation:** Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

### **BOOM LENGTH**

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

### **APPLICATION HEIGHT**

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **SWATH ADJUSTMENT**

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

### **WIND**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **TEMPERATURE AND HUMIDITY**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

## TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small-suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**Spray Volume:** Apply **Tebu-Crop 3.6F** with ground or aerial equipment using sufficient volume of spray to provide thorough coverage. Apply **Tebu-Crop 3.6F** in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Continuous agitation is required to keep the material in suspension.

Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage. Use the higher rate under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.

**Mixing:** Add specified amount of **Tebu-Crop 3.6F** into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the **Tebu-Crop 3.6F** should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. **It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**

**Compatibility:** To determine the compatibility of **Tebu-Crop 3.6F** with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

**Spray Drift Management:** For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wingspan or rotor diameter. Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided. Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

**Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.** Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature. Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**ROTATIONAL CROP RESTRICTIONS:** Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

**Comments:** For optimum disease control, tank mix the lowest specified rate of a spray surfactant with **Tebu-Crop 3.6F**. **Tebu-Crop 3.6F** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Tebu-Crop 3.6F** will be resistant to weathering. **Tebu-Crop 3.6F** is a demethylation inhibitor (DMI) fungicide (Group 3).

| CROP   | DISEASE                      | TEBU-CROP 3.6F APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|--|------------------------------|---|
| Asparagus  | Rust ( <i>Puccinia</i> spp.) | 4 - 6   |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of <b>Tebu-Crop 3.6F</b> (0.11 to 0.17 lb. of active ingredient per acre) in alternation with another effective. Under conditions of severe rust pressure, use the higher rate within the specified rate range. Repeat applications on a 14-day interval as necessary to maintain control of rust.</p> <p><b>Comments:</b> Make applications using ground or aerial application equipment. For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3). Alternating <b>Tebu-Crop 3.6F</b> with other DMI fungicides may lead to resistance.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"><li>• Do not apply to harvestable spears.</li><li>• Do not make more than three foliar applications per season (18 fl. oz./acre or 0.51 lb. of active ingredient per acre).</li><li>• A 50 foot spray drift buffer zone is required for all aerial applications.</li></ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/><b>Pre-harvest interval (PHI) = 100 days (California); 180 days (all other states)</b></p> |                              |   |

| CROP  | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|--|--|
| <b>Barley</b>   | Rusts ( <i>Puccinia</i> spp.)<br>Head Blight ( <i>Fusarium</i> spp.) - Suppression | 4  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. Observe barley fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p><b>Application Timing Directions:</b></p> <p><b>Rusts:</b> Apply <b>Tebu-Crop 3.6F</b> at the earliest sign of rust pustules on foliage.</p> <p><b>Fusarium head blight:</b> Optimal timing of <b>Tebu-Crop 3.6F</b> for fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Apply a maximum of 4 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per crop per season.</li> <li>• Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding.</li> <li>• Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of <b>Tebu-Crop 3.6F</b>.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b></p> |  |  |

| CROP  | DISEASE                                 | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| <b>Beans</b> (fresh & dry, except succulent shelled)  | Rust ( <i>Uromyces appendiculatus</i> ) | 4 – 6  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>Fresh beans:</b> Do not apply more than <b>24 fl. oz. of Tebu-Crop 3.6F</b> per acre per crop season.</li> <li>• <b>Dry beans:</b> Do not apply more than <b>12 fl. oz. of Tebu-Crop 3.6F</b> per acre per crop season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 7 days (fresh beans); 14 days (dry beans)</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|---|---|--|
| <b>Corn</b> (sweet corn, field corn, field corn grown for seed, and popcorn)  | Rust ( <i>Puccinia</i> spp.)<br>Northern Leaf Blight ( <i>Helminthosporium turcicum</i> )<br>Southern Leaf Blight ( <i>Helminthosporium maydis</i> )<br>Northern Leaf Spot ( <i>Helminthosporium carbonum</i> )<br>Gray Leaf Spot ( <i>Cercospora zeae-maydis</i> ) | 4 – 6  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Apply a maximum of 24 fl. oz. (1.5 pints) of <b>Tebu-Crop 3.6F</b> per acre per crop season.</li> </ul> <p><b>Restricted-entry interval (REI) for sweet corn = 19 days</b><br/> <b>Pre-harvest interval (PHI) for sweet corn = 7 days (ears or forage); 49 days (fodder)</b><br/> <b>Restricted-entry interval (REI) for all corn except sweet corn = 12 hours</b><br/> <b>Pre-harvest interval (PHI) for field, seed or popcorn = 21 days (forage); 36 days (grain or fodder)</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| Cotton  | Southwestern Cotton Rust ( <i>Puccinia cacabata</i> ) | 6 – 8  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 24 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per crop season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b></p> <p><b>Pre-harvest interval (PHI) = 30 days</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|---|---|--|
| <p><b>Cucurbit Vegetables Group</b><br/>           Chayote, Chinese Wax gourd, Citron Melon, Cucumber, Gherkin, Edible Gourd (includes Hyotan, Cucuzza, Hechima, and Chinese Okra), <i>Momordica</i> spp. (includes Balsam Apple, Balsam Pear, Bitter Melon and Chinese Cucumber), Muskmelon (includes, Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Clause Melon and Snake Melon), Pumpkin, Summer Squash (includes Crookneck Squash, Scallop Squash, Straightneck Squash, Vegetable Marrow and Zucchini), Winter Squash (includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash and Spaghetti Squash), Watermelon</p> | <p>Powdery Mildew (<i>Sphaerotheca fuliginea/Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)</p> <p>Gummy Stem Blight - Suppression (<i>Didymella bryoniae</i>) (watermelon, squash, pumpkin, and melons only)</p> | <p>4 – 6</p> <p>8</p>                                    |

(continued)

**Cucurbit Vegetables Group (cont.)**

**Notes:** Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.

**Comments:** For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with **Tebu-Crop 3.6F**. **Tebu-Crop 3.6F** must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Tebu-Crop 3.6F** will be resistant to weathering. **Tebu-Crop 3.6F** is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restrictions:**

- Do not apply more than 24 fl. oz. of **Tebu-Crop 3.6F** per acre per crop season.

**Restricted-entry interval (REI) = 12 hours**

**Pre-harvest interval (PHI) = 7 days**

| CROP   | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE)                                   |
|--|---|--|
| Dry Bulb Onion<br>Garlic, Great-headed (Elephant)  | White Rot ( <i>Sclerotium cepivorum</i> )   | White Rot: 20.5 fl. oz. per acre applied in a 4 to 6 inch band over/into each furrow |
| Garlic<br>Shallot  | Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> )<br>Purple Blotch ( <i>Alternaria porri</i> ) | 4 – 6  |
| <p><b>White Rot:</b> For the control of white rot, make one application in the furrow at the time of planting. Make the in-furrow application at the rate of 20.5 fl. oz. <b>Tebu-Crop 3.6F</b> per acre. Apply the entire per acre rate in a 4- to 6-inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 – 6 fl. oz./acre.</p> <p><b>Rust:</b> For the control of rust, make foliar applications at the rate of 4 – 6 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per application. Repeat at an interval of 10 – 14 days. Apply <b>Tebu-Crop 3.6F</b> in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Comments:</b> For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 32.5 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season if an in-furrow treatment is made. If <b>Tebu-Crop 3.6F</b> is not applied as an in-furrow treatment, then do not apply more than 12 fl. oz./acre per season as a foliar spray.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 7 days</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| <p><b>Fruiting Vegetable Group 8-10 (Except Okra)</b><br/> African Eggplant, Bush Tomato, Bell Pepper, Cocona, Currant Tomato, Eggplant, Garden Huckleberry, Goji Berry, Ground Cherry, Martynia, Naranjilla, Pea Eggplant, Pepino, Non-bell Pepper, Roselle, Scarlet Eggplant, Sunberry, Tomatillo, Tomato, Tree Tomato; Cultivars, varieties, and/or hybrids of these.</p>  | <p>Early Blight<br/> (<i>Alternaria solani</i>)</p> | <p>8</p>   |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> as a foliar spray using an interval of 7 days.</p> <p><b>Comments:</b> For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 48 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 7 days</b></p> |   |  |

| CROP   | DISEASE  | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|--|--|--|
| <b>Garden Beet</b><br>roots and tops<br>(leaves)   | <i>Cercospora</i> Leaf Spot ( <i>Cercospora beticola</i> ) | 3 – 7.2  |
| <p><b>Notes:</b> Make applications on 14-day intervals.</p> <p><b>Comments:</b> For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on beet foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 28.8 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b></p> <p><b>Pre-harvest interval (PHI) = 7 days</b></p> |  |  |

| CROP   | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|--|--|--|
| <b>Grasses Grown for Seed</b>  | Rust ( <i>Puccinia</i> spp.)<br>Powdery Mildew | 4 – 8  |
| <p><b>Notes:</b> Apply the specified rate of <b>Tebu-Crop 3.6F</b> in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit the lowest specified rate of a spray surfactant should be tank mixed with <b>Tebu-Crop 3.6F</b>.</p> <p><b>Rust:</b> Apply the specified rate of <b>Tebu-Crop 3.6F</b> as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.</p> <p><b>Powdery Mildew:</b> Apply specified rate of <b>Tebu-Crop 3.6F</b> when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 16 fl. oz. <b>Tebu-Crop 3.6F</b> (.45 lb. a.i.) per acre per crop season.</li> <li>• Chaff, screenings and straw from treated areas may be used for feed purposes. Do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</li> </ul> <p><b>Pre-harvest interval (PHI) = 4 days</b><br/> <b>Restricted-entry interval (REI) = 12 hours</b></p> |  |  |

| CROP   | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|--|--|--|
| <b>Green Onion</b><br><b>Leek</b><br><b>Spring Onion</b><br><b>Welsh Onion</b>   | White Rot ( <i>Sclerotium cepivorum</i> )<br>(suppression only)<br>Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> )<br>Purple Blotch ( <i>Alternaria porri</i> ) | 4 – 6  |
| <p><b>Notes:</b> For the control of diseases, make foliar applications using an interval of 10 – 14 days. Apply <b>Tebu-Crop 3.6F</b> in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Comments:</b> For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 24 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 7 days</b></p> |  |  |

| CROP  | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|--|--|
| Hops  | Powdery Mildew ( <i>Sphaerotheca humuli</i> /<br><i>Sphaerotheca macularis</i> ) | 4 – 8  |
| <p><b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Increase the spray volume and the application rate as vine growth increases during the season.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 32 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per crop season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 14 days</b></p> |  |  |

| CROP   | DISEASE   | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|--|---|--|
| <b>Leafy Brassica Greens</b><br>Broccoli Raab<br>Chinese Cabbage (Bok Choy)<br>Collards<br>Kale<br>Mizuma<br>Mustard Greens<br>Mustard Spinach<br>Rape Greens<br>Turnip Greens   | Cercospora Leaf Spot<br><i>(Cercospora brassicicola)</i><br>Powdery Mildew ( <i>Erysiphe cruciferarum</i> )<br>Alternaria Leaf Spot<br><i>(Alternaria brassicicola)</i> | 3 – 4  |
| <p><b>Comments:</b> For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 16 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> <li>• Application to turnip greens is limited to east of the Rockies.</li> </ul> <p><b>Reapplication interval: Do not apply more than once every 10 days.</b></p> <p><b>Restricted-entry interval (REI) = 12 hours</b></p> <p><b>Pre-harvest interval (PHI) = 7 days</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| Lychee  | Anthracnose ( <i>Colletotrichum gloeosporioides</i> ) | 4 – 6  |
| <p><b>Notes:</b> Begin first application of <b>Tebu-Crop 3.6F</b> as panicle emerges. Spray up to 6 fl. oz. <b>Tebu-Crop 3.6F</b> per acre every 10 days thereafter for a total of 8 sprayings. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a nonionic spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 48 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 2 days</b><br/> <b>Pre-harvest interval (PHI) = 0 (zero) days</b></p> |   |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| Okra  | <i>Cercospora</i> Leaf Spot ( <i>Cercospora</i> spp.) | 4 – 6  |
| <p><b>Notes:</b> Apply specific dosage of <b>Tebu-Crop 3.6F</b> in a preventative spray program. Use the highest specified rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 24 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 3 days</b></p> |   |  |

| CROP  | DISEASE  | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|---|--|--|
| Peanut  | <b>SOILBORNE:</b><br>Cylindrocladium Black Rot (Suppression)<br>Rhizoctonia Limb Rot<br>Rhizoctonia Pod Rot (Virginia and North Carolina only)<br>Sclerotium Stem and Pod Rot (White Mold,<br>Southern Blight Southern Stem Rot) | 7.2  |
|   | <b>FOLIAR:</b><br>Early Leaf Spot<br>Late Leaf Spot<br>Leaf Rust<br>Pepper Spot ( <i>Leptosphaerulina</i> )<br>Web Blotch ( <i>Phoma</i> )   | 7.2  |
| <p><b>Notes:</b> For optimum control of the specified soilborne diseases, four consecutive applications of <b>Tebu-Crop 3.6F</b> must be made at 14-day intervals.</p> <p><b>Tebu-Crop 3.6F</b> is a sterol demethylation inhibitor (DMI) fungicide. Tank mix Chlorothalonil at the rate of 12 ounces of chlorothalonil active ingredient with <b>Tebu-Crop 3.6F</b> at the specified rate as a leaf spot resistance management strategy. A spray surfactant is not necessary when <b>Tebu-Crop 3.6F</b> is tank mixed with chlorothalonil. Mixing or alternating <b>Tebu-Crop 3.6F</b> with other DMI fungicides may lead to resistance.</p> <p><b>Tebu-Crop 3.6F</b> must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of <b>Tebu-Crop 3.6F</b> against the root and pod rots. Use <b>Tebu-Crop 3.6F</b> in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.</p> <p style="text-align: right;"><i>(continued)</i></p> |  |  |

### **Peanut (cont.)**

**FOUR-APPLICATION SPRAY PROGRAM:** Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Make applications of chlorothalonil prior to and following applications of **Tebu-Crop 3.6F** to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, tank mix the lowest specified label rate of a spray surfactant with **Tebu-Crop 3.6F**.

**LEAF SPOT ADVISORY SCHEDULE:** For control of soil-borne diseases in an advisory schedule, apply **Tebu-Crop 3.6F** in the first advisory spray in July and continue **Tebu-Crop 3.6F** applications at 14-day intervals. When applying **Tebu-Crop 3.6F** after August 15, tank mix with chlorothalonil for resistance management purposes.

#### **Application Timing of Tebu-Crop 3.6F for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot**

| <b>Spray Program</b> | <b>Tebu-Crop 3.6FL Fungicide Application No.</b> | <b>Chlorothalonil Application No.</b> |
|----------------------|--|---------------------------------------|
| applications         | 3, 4, 5 and 6                                    | 1, 2 and 7                            |

#### **Restrictions:**

- Apply a maximum of 28.8 fl. oz. (.81 lb. a.i.) of **Tebu-Crop 3.6F** per crop season.
- Do not feed hay or threshings or allow livestock to graze in treated areas.

**Restricted-entry interval (REI) = 12 hours**

**Pre-harvest interval (PHI) = 14 days**

| CROP   | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|--|--|--|
| <b>Pecan</b>   | Brown Leaf Spot ( <i>Sirosporium diffusum</i> )<br>Downy Spot ( <i>Mycosphaerella caryigena</i> )<br>Liver Spot ( <i>Gnomonia caryae</i> )<br>Scab ( <i>Cladosporium caryigenum</i> )<br>Vein Spot ( <i>Gnomonia nerviseda</i> )<br>Zonate Leaf Spot ( <i>Grovesinia pyramidalis</i> ) | 4 – 8  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> in a preventative spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Apply <b>Tebu-Crop 3.6F</b> 4 fl. oz./acre in a tank-mix with the specified rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin®. Do not add a surfactant to the spray solution when tank-mixing <b>Tebu-Crop 3.6F</b> with Super-Tin®. Apply <b>Tebu-Crop 3.6F</b> in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 – 8 fl. oz./acre of <b>Tebu-Crop 3.6F</b> to full-size mature trees, and 4 – 6 fl. oz. <b>Tebu-Crop 3.6F</b> per acre to smaller trees. Apply the higher rate within the specified rate range to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. Add the lowest specified rate of a surfactant to the spray solution for optimum control of the indicated diseases.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3). Apply in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply after shucks begin to split. Do not apply more than 32 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per crop season.</li> <li>• Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = Do not apply after shucks begin to split.</b></p> |  |  |

| CROP  | DISEASE   | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|---|--|
| Soybean   | Rust ( <i>Phakopsora pachyrhizi</i> )<br>Powdery Mildew ( <i>Microsphaera diffusa</i> ) | 3 – 4  |
| <p><b>Notes:</b> Apply <b>Tebu-Crop 3.6F</b> as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher specified rates within the specified rate range and use shorter specified spray intervals when disease pressure is severe. The lowest specified label rate within the specified rate range of a spray surfactant may be tank mixed with <b>Tebu-Crop 3.6F</b>. Apply <b>Tebu-Crop 3.6F</b> in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 12 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per use season.</li> <li>• Do not make more than three applications per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 21 days</b></p> |   |  |

| CROP   | DISEASE                            | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|--|------------------------------------|--|
| <b>Sunflower</b>   | Rust ( <i>Puccinia helianthi</i> ) | 4 – 6  |
| <p><b>Notes:</b> Apply specific dosage of <b>Tebu-Crop 3.6F</b> at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply the higher specified rate within the specified rate range to highly susceptible varieties and/or under severe disease conditions. Repeat application at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. Contact your State extension service for a list of approved surfactants. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 16 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 50 days</b></p> |                                    |  |

| CROP   | DISEASE  | TEBU-CROP 3.6F<br>APPLICATION RATE<br>(FL. OZ. PER ACRE) |
|--|--|--|
| <b>Turnip</b> (Application is limited to east of the Rockies)  | Cercospora Leaf Spot<br>( <i>Cercospora brassicicola</i> ) | 4 – 7.2  |
| <p><b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals.</p> <p><b>Comments:</b> For optimum disease control, tank mix the lowest specified rate within the specified rate range of a spray surfactant with <b>Tebu-Crop 3.6F</b>. <b>Tebu-Crop 3.6F</b> must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, <b>Tebu-Crop 3.6F</b> will be resistant to weathering. <b>Tebu-Crop 3.6F</b> is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 28.8 fl. oz. <b>Tebu-Crop 3.6F</b> per acre per crop season.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b><br/> <b>Pre-harvest interval (PHI) = 7 days</b></p> |  |  |

| CROP  | DISEASE  | TEBU-CROP 3.6F APPLICATION RATE (FL. OZ. PER ACRE) |
|---|--|--|
| Wheat   | Rusts: leaf, stem, and stripe<br>( <i>Puccinia</i> spp.)<br><br>Head blight or scab<br>( <i>Fusarium</i> spp.) - Suppression | 4  |
| <p><b>Notes:</b> Observe wheat fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Apply <b>Tebu-Crop 3.6F</b> in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><b>Application Timing Directions:</b></p> <p><b>Rusts:</b> Apply <b>Tebu-Crop 3.6F</b> at the earliest sign of rust pustules on foliage.</p> <p><b>Fusarium head blight:</b> Optimal timing of <b>Tebu-Crop 3.6F</b> for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Apply a maximum of 4 fl. oz. of <b>Tebu-Crop 3.6F</b> per acre per crop per season.</li> <li>• Straw cut after harvest may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with <b>Tebu-Crop 3.6F</b>.</li> </ul> <p><b>Restricted-entry interval (REI) = 12 hours</b></p> <p><b>Pre-harvest interval (PHI) = 30 days</b></p> |  |  |

### **SEED TREATMENT – Corn** (Sweet Corn, Field Corn Grown for Seed, and Popcorn)

For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

#### **Seed Bag Label Requirements**

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with **Tebu-Crop 3.6F**, a fungicide containing tebuconazole.
- Do not use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

**USE RESTRICTION:** When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

| DISEASE  | RATE Fl. Oz./<br>CWT | DIRECTIONS FOR USE   |
|--|----------------------|--|
| <b>Soilborne and Seedborne</b><br>Fusarium                                       | 0.071                | Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Dilute product with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates specified for the crop to be treated with <b>Tebu-Crop 3.6F</b> . The length of control will vary depending on the rate used. |
| <b>Soilborne and Seedborne</b><br>Head Smut<br>( <i>Sphacelotheca reiliana</i> ) | 0.27 – 0.54          |  |

### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in the original container in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed. Store out of the reach of children, preferably in a locked storage area. Open and handle container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Nonrefillable container. Do not refill or reuse container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate in application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC, Manufacturer and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC AND MANUFACTURER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL To the extent consistent with applicable law, Sharda USA LLC, Manufacturer or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC, MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC, MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Sharda USA LLC, Manufacturer and Seller offer this product, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

# Tebu-Crop 3.6F

**ACTIVE INGREDIENT:** % BY WT.

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-  
alpha(1,1-dimethylethyl)-1H-1,2,4-  
triazole-1-ethanol ..... 38.7%

**OTHER INGREDIENTS:** ..... 61.3%

**TOTAL:** ..... **100.0%**

Contains 3.6 pounds tebuconazole per gallon

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

See complete Precautionary Statements, and Directions For Use, including Storage and Disposal in booklet.

**STOP-Read the label before use.**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Manufactured For:** Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707  
**EPA Reg. No. 83529-11 EPA Est. No. 70815-GA-001 Net Contents: 2.5 Gallons**

### FIRST AID

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **If On Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically. **Symptoms of Poisoning:** The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

**EMERGENCY NUMBERS:** For 24-hour medical emergency assistance (human or animal) call **1-800-222-1222**. For chemical emergency assistance (spill, leak, fire, or accident) call CHEMTREC at **1-800-424-9300**.